

AMPROL[®] Vet Powder

Effective for the prevention and treatment of all types of Coccidiosis in Poultry

Composition

Each gram powder contains:

Amprolium hydrochloride BP 200 mg

Mode of action

As the structure of **Amprol[®]** Vet powder resembles to that of Vitamin B1, it acts as antagonist of Vitamin B1. As a result, due to the lack of essential Vitamin B1, protozoa cannot reproduce and ultimately destroyed.

Indications

Amprol[®] Vet powder is used in the following diseases of chicken, broiler and layer poultry, such as; Coccidiosis or Blood dysentery caused by *Eimeriaacervulina* (duodenum), *E. maxima* and *E. necatrix* (small intestine), *E. brunetti* (large intestine) and *E. tenella* (caecum).

Dosage & Administration

Poultry

For Prevention:

Water: 30 gm / 100 liters of drinking water for 7-14 consecutive days.

Feed: 60-75 gm / 100 kg of feed for 7- 14 consecutive days.

For Treatment:

Normal case: 60 gm / 100 liters of drinking water for 5-7 consecutive days.

In severe case: 120 gm / 100 liters of drinking water for 5-7 consecutive days.

Calves/lamb

For prevention: 1 gm / 40 kg of body weight for 5 consecutive days.

For treatment: 2 gm / 40 kg of body weight for 5 consecutive days.

Or, as per direction of registered veterinarian / Consultant.

Contraindications

Amprol[®] Vet powder should not be used in combination with Thiamin (Vitamin B1).

Withdrawal period

Meat: 0 day;

Egg: 0 day

Precautions

Store in a dry place, away from light. Keep out of the reach of children. Once opened the sachet should be used quickly.

Side effect

There is no side effect for recommended dosage.

Supply

100g & 500g

For Veterinary Use Only