

Verkil™ Vet

Ivermectin BP 1% Oral Solution

Composition

Each ml **Verkil™ Vet** solution contains Ivermectin BP 10mg.

Indications

Verkil™ Vet acts against following endo and ecto parasites:

Endo-parasites: Round worm

Ecto-parasites: Tick, Lice, Mite, Flea

Pharmacology

Verkil™ Vet have strong affinity to glutamate gated chloride ion channel of nerves and causes hyper-polarization of the nerve or muscles cells and resulting paralysis and killing of parasites. Ivermectin is metabolized in the liver and its metabolites are excreted almost exclusively in the feces over an estimated 12 days with less than 1% of the administered dose excreted in the urine.

Dosage & Administration

Livestock: Pour on slowly from the withers to tail head at a dose of 1 mL/20 kg body weight.

Poultry: Oral dose:

Endo-parasites: 1 mL/ 25 kg body weight (Ivermectin 0.4 mg/kg body weight) for 1 day. if necessary repeat the dose after 15 days

Ecto-parasites: 1 mL/ 25 kg body weight (Ivermectin 0.4 mg/kg body weight) for 2 days. if necessary repeat the dose after 15 days.

Pond: 1mL/ decimal (per feet depth of water)

Or, as per direction of registered Veterinarian/Consultant.

Contraindications

Not known.

Precautions

Once opened the pack should be used quickly. Store in a cool and dry place, away from light. Keep out of the reach of children.

Side effect

Not known.

Supply

15 mL, 100 mL & 500 mL

For Veterinary / Aquaculture Use Only