

Povidon Vet[®]

Povidon Iodine 10% USP Solution

Composition

Povidon Iodine 10% USP Solution

Indication

- Povidon Vet Controls Viral, bacterial, fungal diseases and certain protozoan in shrimp/Fish, Poultry & Dairy
- Povidon Vet deodorize and purifies pond water
- Povidon vet Used as Disinfectant & to maintain Bio-security in Aquaculture, poultry & Dairy Farm

Dosage & Administration

Aqua:

Hatchery fry & eggs disinfection: 2-3 mL/100 liter water (10 min dip bath)

Brood fish disinfection: 3-5 mL/100 liter water (10 min dip bath)

Pond or Gher: 1-1.5 liter/ Acre for 3-4 ft. depth

Poultry:

Poultry Farm, utensils, equipment's & footbath: 7 mL/liter water

Cattle:

Udder disinfection: 4 mL/liter water

Water disinfection: 6-8 mL/liter water

Direction of use

Take the required quantity of Povidon vet and mix it with pond water, apply the mix throughout the pond. In poultry/cattle mix povidon vet with water and spray/apply in required site. Or as direction by Aquaculture/ Veterinary consultant

Contraindications

Not known.

Precautions

Once opened the pack should be used quickly. Store in a cool and dry place, away from light. Keep out of the reach of children

Side effect

Not known.

Supply

100 mL, 500 mL & 1 L

For Aquaculture/Poultry Use Only