

Heatler® -VET

Electrolytes and Betaine oral solution

Composition

Each ml contains

Analyzed

Betaine Anhydrous	250.00 mg
Magnesium Chloride	18.75 mg
Potassium Chloride	49.45 mg
Sodium Chloride	22.75 mg
Sodium Metabisulfite	50.33 mg

Enriched

Glycine, Dextrose, Anise Oil, Propylene glycol, Inositol, Polyethylene Glycol 400, Polysorbate 20, Citric acid.

Usage

- To maintain electrolyte balance during diarrhea and dehydration
- To prevent Heat Stress
- Before and after transportation
- During and after antibiotic treatment and Vaccination
- To increase water intake and water holding capacity of the animal body

Dosage & administration

Species	Condition	Dose
Poultry	Dehydration, Diarrhea, Electrolyte imbalance or Normal condition	1-2 ml/10L drinking water for 5-10 days.
	Heat Stress	1.5-2.5 ml/10L drinking water for 5-10 days.
	Transportation Associated Stress <i>Dose should be administered both before and after transportation but not longer than 48 hours in a row</i>	Before Transportation: 8ml/10L drinking water. The dose should be administered 24 hours before transportation. After Transportation: 8ml/10L drinking water. The dose should be administered right after transportation.
Cattle	Dehydration, Diarrhea, Electrolyte imbalance	Dissolve in 25 ml/L water, Administer twice a day for 1-6 days.

Contraindication

Not known

Precaution

Store in cool and dry place, away from light. Keep out of the reach of children. Once opened the container should be used quickly.

Side effect

Not known.

Supply

100 mL & 500 mL Bottle.

Animal Feed Premix