

PROTIMIN[®] SOLUTION

Combination of 18 Amino acids & 3 minerals oral solution

Composition

Each 100ml solution contains

L-Lysine (As hydrochloride) USP	1250.00 mg
L-Methionine USP	500.00 mg
L-Threonine USP	620.00 mg
L-Tryptophan USP	90.00 mg
Glycine USP	760.00 mg
L-Histidine USP	250.00 mg
L-Arginine USP	500.00 mg
L-Alanine USP	200.00 mg
L-Aspartic acid USP	250.00 mg
L-Cystine USP	10.00 mg
L-Glutamic acid USP	75.00 mg
L-Isoleucine USP	352.00 mg
L-Leucine USP	490.00 mg
L-Phenylalanine USP	533.00 mg
L-Proline USP	100.00 mg
L-Serine USP	100.00 mg
T-Tyrosine USP	25.00 mg
L-Valine USP	360.00 mg
Manganese (As Manganese Sulfate M/H)	1.6 mg
Iron (As Dried Ferrous Sulfate)	4.00 mg
Zinc (As Zinc Sulfate)	2.00 mg

Usages

Protimin[®] solution can be used in deficiency of amino acids and minerals. **Protimin[®]** solution can also be used in the following conditions:

- ❖ To increase body weight and feathering
- ❖ To reduce stress due to food change, transportation, debaking, temperature change.
- ❖ To improve the effect of vaccination
- ❖ To increase immunity
- ❖ Reduce the stress due to antibiotic and anthelmintic
- ❖ To reduce cannibalism
- ❖ To increase fertility & hatchability of egg
- ❖ To increase the size & production & laying period of egg
- ❖ To increase the laying period

Dosage & administration

Poultry: 1ml/1-2Liter of drinking water for 5-7 days

Cattle: 10-20ml per animal daily

Or, as per direction of registered veterinarian / Consultant.

Contraindication

Not found.

Precaution

Store in a dry place, away from light. Keep out of the reach of children.

Withdrawal period

Egg: 0 (zero) day

Meat: 0 (zero) day

Side effect

Not known.

Supply

100 ml, 500 ml and 1 L bottle

Animal / Poultry Feed Premix